

Pulmonary Function after Adenotonsillectomy

Mehrdad Rogha^{1,*}, Jaleh Amini¹, Mostafa Raisi²

Abstract

Introduction

Adenotonsillar hypertrophy is a common disorder among children which, without proper treatment, may lead to considerable problems. Although the consequences of this disorder have been studied in other articles, we decided to evaluate the changes in pulmonary function tests in these children after adenotonsillectomy, and the correlation between clinical and spirometric parameters.

Materials and Methods:

We conducted a before- and after- clinical trial. Forty children (17 females and 23 males) with a diagnosis of upper airway obstruction due to adenotonsillar hypertrophy were enrolled in this study. Mean age of the participants was 6.9 ± 1.9 years. Eight spirometric parameters were selected for evaluation pre-operatively and 40 days postoperatively. Besides, symptom scores were defined for each patient to assess their disease severity, pre- and postoperatively. Data were analyzed statistically.

Results:

Forced vital capacity (FVC) increased from $1.28 \pm 0.26\%$ pre-operatively to $1.33 \pm 0.24\%$ postoperatively ($P=0.05$). Peak expiratory flow increased from $2.74 \pm 0.65\%$ pre-operatively to $2.84 \pm 0.51\%$ postoperatively ($P=0.02$) and mid expiratory forced expiratory flow (FEF_{25-75}) was $1.81 \pm 0.48\%$ pre-operatively, increasing to $1.91 \pm 0.50\%$ postoperatively ($P=0.02$). Maximal expiratory flow at 25% of FVC (MEF_{25}) increased from $1.09 \pm 0.36\%$ pre-operatively to $1.21 \pm 0.34\%$ postoperatively ($P=0.02$). There was no correlation among the other spirometric parameters (FEV_1 , FEV_1/FVC , MEF_{50} and MEF_{75}) pre- and post-operatively ($P>0.05$). Despite some improvements in pulmonary function indices, there was no correlation between changes in spirometric parameters and severity of the snoring ($P>0.05$).

Conclusion:

Although our findings reveal that adenotonsillectomy had a positive effect on pulmonary function tests, we found no significant correlation between alterations in spirometric parameters and severity of snoring. However, performing a spirometric examination in children with adenotonsillar hypertrophy may be beneficial for assessing the pulmonary status of the affected patient.

Keywords:

Adenotonsillar hypertrophy, Adenoidectomy, Pulmonary function test, Upper airway obstruction.

Received date: 16 May 2016

Accepted date: 2 Oct 2016

¹Department of Otorhinolaryngology, Isfahan University of Medical Science, Isfahan. Iran.

²Poursina Hakim Research Institute, Isfahan University of Medical Science, Isfahan, Iran.

*Corresponder Author:

Department of Otorhinolaryngology, Isfahan University of Medical Science, Isfahan. Iran.

Tel: 09133087832, E-mail: ja61057@yahoo.com

Introduction

Adenotonsillar hypertrophy (ATH) is a common disorder in children (1,2), and the most common cause of upper airway obstruction in this age group (3). The role of ATH as an important etiology in chronic airway obstruction and ventilatory impairment during sleep has been established (4,5). Without the proper treatment, ATH might lead to considerable problems, such as behavioral, cognitive, and cardiovascular concerns (6-10). Disturbed inspiration in ATH may present with hypoxia, hypercarbia, and elevated bicarbonate levels (11), as well as upper airway obstruction and resultant mouth breathing, often leading to pulmonary morbidities (12).

Fortunately, adenotonsillectomy results in a noticeable improvement in quality of life, cognition, and behavior of patients (13,14), but it is still unclear whether it can affect pulmonary function. Till now, few studies have evaluated the effect of ATH and adenotonsillectomy on pulmonary function tests (PFT).

PFTs in children with ATH were examined by Maurizi et al, who concluded that 52% of patients showed parameters indicating lower airway obstruction (15). An experimental study reported by Di Martino et al. revealed that adenoids may have an impact on expiratory flow (16).

Identification of upper airway obstruction using a peak flowmeter and spirometer has been proposed by some experts, with significant changes between pre- and post-surgery spirometric parameters reported (3,12,15,17). Most investigators have suggested that changes in pulmonary function are transient and disappear after treatment (18). Based on our research, no studies have been published on the correlation between clinical and spirometric parameters.

Thus, we conducted this study in order to evaluate PFT in children affected by symptomatic upper airway obstruction due to ATH, and compare pre-and post-operative values to evaluate any correlation between clinical and spirometric parameters.

Materials and Methods

In a prospective trial conducted from September 2015 to April 2016 in the department of Otorhinolaryngology and Pulmonology of AL Zahra Hospital, Isfahan, Iran, 40 boys and girls

(5–14 years of age) were recruited from patients waiting for an adenotonsillectomy procedure. All cases presented with enlarged tonsils graded >2+ (degrees III and IV), according to the Brodsky classification and a lateral head X-ray proving significant adenoid hypertrophy with an adenoid-nasopharyngeal (A/N) ratio >0.70 according to the Fujioka method (19,20). Cases excluded from the group were those with cardio-pulmonary disease, neurological involvement, obese children (body mass index [BMI] >30kg/m²), those with no surgical indication, and those with other causes of nasal obstruction such as polyps, nasal septal deviations, and thoracic skeletal deformity, for example.

Spirometry is a simple non-invasive method that evaluates only volumes above the residual volume and flow volume rates (21). Spirometric values were measured using a spirometer (P.K. Morgan Ltd., UK). Spirometric values were obtained with the patients seated, with clipped nose and with no restrictions to chest expansion, such as fitted or tight clothes, and no orthodontic braces. To obtain accurate results, cases were encouraged to attempt the test a maximum of five times and the maximal result was recorded. All measurements were completed during a maximum 2-minute interval.

The respiratory test was carried out by the same operator, after careful training. The operator and gave plenty of encouragement to the children to perform the tests.

Children were assessed twice; 1 week before and 40 days after surgery in the same clinic and by the same operator. As well as spirometric parameters, snoring and duration of disease and its associated symptoms such as enuresis, open mouth breathing, and hyponasal sound were evaluated through questionnaires pre- and post-operatively. The severity of snoring was evaluated using a visual analog scale (VAS). The significance of changes in respiratory indices and the correlation between spirometric values and symptom scores were analyzed by Student's and McNemar's tests.

The protocol was approved by the Ethics and Research Committee of Isfahan University of Medical Sciences. Parental consent was also signed before evaluation for each patient.

Results

The sample group consisted of 42 children (19 female and 23 male) aged between 5 and 11

Pulmonary Function after Adenotonsillectomy

years (mean, 6.9±1.7 years). Two female children were excluded from the study because relocation meant that they did not perform the second procedure. Therefore, the analysis was performed on 40 cases. All patients presented with nasopharyngeal obstruction (A/N ratio > 0.70). Twenty-nine patients exhibited grade III

and the others had the grade IV enlarged tonsils. Table 1 shows the improvement in forced vital capacity (FVC), peak expiratory flow (PEF), maximal expiratory flow at 25% of FVC (MEF₂₅), and mid expiratory forced expiratory flow (FEF₂₅₋₇₅) 40 days after surgery as compared with preoperative measures (P=0.02).

Table 1: Mean Pre-adenotonsillectomy and Post-adenotonsillectomy spirometric parameters.

Spirometric Parameters	Pre-adenotonsillectomy (Mean±SD)	Post-adenotonsillectomy (Mean±SD)	P-value
FEV ₁	1.21±0.23	1.23±0.21	0.41
FVC	1.28±0.26	1.33±0.24	0.04
FEV ₁ /FVC	95±4	93±5	0.07
FEF ₂₅₋₇₅	1.81±0.5	1.91±0.48	0.02
PEF	2.74±0.65	2.84±0.51	0.02
MEF ₂₅	1.09±0.36	1.21±0.34	0.02
MEF ₅₀	2.13±0.47	2.09±0.50	0.384
MEF ₇₅	2.68±0.57	2.72±0.45	0.498

SD: standard deviation, FEV₁: forced expiratory volume during the first second of expiration, FVC: forced vital capacity, FEV₁/FVC: forced expiratory volume during the first second of expiration to forced vital capacity, FEF₂₅₋₇₅: midexpiratory forced expiratory flow, MEF₂₅: maximal expiratory flow at 25% of FVC, MEF₅₀: maximal expiratory flow at 50% of FVC, MEF₇₅: maximal expiratory flow at 75% of FVC.

We found no statistically significant differences in forced expiratory volume during the first second of expiration (FEV₁), forced expiratory volume during the first second to forced vital capacity (FEV₁/FVC) ratio, maximal expiratory flow at 50% of FVC (MEF₅₀), or maximal expiratory flow at 75% of FVC (MEF₇₅) for the same period (P>0.05).

Snoring score was obtained using a VAS. The mean VAS score decreased from 62.9±30.9 pre-operatively to 2.9±4.6 post-operatively (P<0.001).

Before adenotonsillectomy, 11 patients (27.5%) reported enuresis, 34 patients (85%) experienced hyponasal sound, and 34 patients (85%) reported mouth breathing. Forty days after the operation, these numbers changed to three (7.5%), two (5%), and two (5%), respectively (Fig.1).




Fig 1: frequency of enuresis, Hyponasal sound and mouth breathing pre and post adenotonsillectomy

There was no correlation between changes in spirometric parameters and snoring and the changes assessed in terms of VAS (Table.2). Changes in FEF₂₅₋₇₅ were correlated with changes in other spirometric parameters (P<0.05, r>0); i.e. improvement in each of the spirometric parameters was concomitant with improvement in FEF₂₅₋₇₅ and vice versa.

Table 2: Pearson's correlation coefficient in changed spirometric parameters and snoring and its changes.

Changed spirometric parameters	VAS of snoring		VAS of changes in snoring	
	r	P-value	r	P-value
FEV ₁	-0.15	0.17	0.12	0.23
FVC	-0.14	0.19	0.12	0.23
FEV ₁ /FVC	0.04	0.40	0.02	0.44
FEF ₂₅₋₇₅	-0.98	0.27	0.08	0.32
PEF	0.09	0.29	-0.11	0.25
MEF ₂₅	-0.19	0.11	0.18	0.14
MEF ₅₀	-0.16	0.16	0.12	0.23
MEF ₇₅	-0.06	0.35	0.03	0.43

VAS: visual analogue scale, r: Pearson's correlation coefficient, FEV₁: forced expiratory volume during the first second of expiration, FVC: forced vital capacity, FEV₁/FVC: forced expiratory volume during the first second of expiration to forced vital capacity, FEF₂₅₋₇₅: midexpiratory forced expiratory flow, MEF₂₅: maximal expiratory flow at 25% of FVC, MEF₅₀: maximal expiratory flow at 50% of FVC, MEF₇₅: maximal expiratory flow at 75% of FVC.

Discussion

Upper airway obstruction can be evaluated by a non-invasive technique such as PFT (spirometry) which helps to detect those obstructive changes that might not be clinically evident (22).

In the present study, we found improvement in FVC, PEF, MEF_{25} and FEF_{25-75} 40 days after surgery in children with enlarged tonsils as compared with baseline measures. This finding is in contrast with a number of other studies (12,15,23). For example, Yadav et al. reported significant improvement in FIF_{50} and FEF_{50}/FIF_{50} , $FEV_1/PEFR$, and $FEV_1/FEV_{0.5}$ ratios following surgery (22). Similar to our findings, Kavukcu et al. reported an increase in FVC, PEF, FEV_1 , MEF_{25} , FEF, and FEF_{75} after adenotonsillectomy (18).

Some authors believe that the lower airway shows obstructive changes in the children with ATH. They believe that disorders of the upper and lower airways commonly coexist, as they present a similar histology that may be considered in the context of united airways (24,25).

On the other hand, obstruction of the upper airway can cause alterations in other parts of the respiratory system, as well as in the amount of muscle stretch and thoracic movement (26). Children with ATH tend to have trouble with nasal breathing, and therefore prefer to breathe through the mouth (12). It has been also assumed that less nasal humidification and warming due to mouth breathing can lead to alterations in the diffusion and viscosity of the surfactant which may be a powerful stimulus for bronchiolar obstruction (27). Chronic mouth breathing may also impose some undesired physical effects on respiratory function, by deforming the craniofacial skeleton, oral cavity and posture, and also on psychological characteristics (28).

Moreover, relative hypoxemia and hypercarbia are seen in many patients with ATH (11). The decrease in both maximum breathing capacity and respiratory response to CO_2 (29) may be responsible for alterations in pulmonary function, as seen in the functional indices of spirometry. Furthermore, tonsils and adenoids are the principal parts of Waldeyer's ring which its basic role is antibody synthesis. Chronic ATH is recognized by chronic inflammation and hypoxia/reoxygenation episodes. This may lead to synthesis of inflammatory cytokines, and free nitrogen and oxygen radicals (30, 31) which

may be responsible for impaired pulmonary function.

Finally, as described by other researchers, inflammation of the nasal and oropharyngeal mucosa is a common finding, as in adults with sleep-disordered breathing (SDB) (32). C-reactive protein, a systemic indicator for inflammation, has recently been found to be elevated in the serum of children with ATH (33). Additionally, expression of glucocorticoid receptors in the upper airway lymphoid tissues and expression of inflammatory mediators and leukotriene receptors were increased in children with SDB (34-36).

We suggest that the increased inflammatory conditions have some effect on the upper and lower airway in these children. The mechanisms which promote this inflammatory reaction are probably related, at least partially, to the vibratory mechanical injury associated with snoring (37). However, we found no correlation between clinical findings such as snoring and changes in spirometric parameters.

In our study, we observed changes in a number of obstructive spirometric indices. Our study was designed to be as simple as possible with a low cost. This simplicity facilitates the reproducibility of similar studies in routine medical practice. On the other hand, it is often difficult to obtain accurate measurements through spirometry because of poor cooperation in performing the test in children (18). PFT could reveal the obstructive influence of ATH and could be valuable in surgical indications of ATH due to recurrent infections in children (18,22). It could also be beneficial in convincing hesitant parents of the importance of a timely decision with respect to adenotonsillectomy.

Conclusion

Although adenotonsillectomy might have a positive effect on the function of the lower respiratory tract, we found no significant correlation between changes in spirometric parameters and snoring. However, performing a spirometric examination in children with ATH may be beneficial for assessing the pulmonary function of these patients.

It seems that PFT can reveal the obstructive effect of ATH whenever slight clinical or radiological obstructive signs are present, and might be valuable in surgical indications of ATH due to recurrent tonsillar infections in

children with a mildly obstructed airway. Further studies are needed to explore the precise etiology of these changes.

Acknowledgments

This manuscript is supported by grants from the Isfahan University of Medical Sciences. The authors acknowledge the work of Dr. Masoud Maleki in designing the study and in carrying out data collection.

References

1. Khalifa MS, Kamel RH, Zikry MA, Kandil TM. Effect of enlarged adenoids on arterial blood gases in children. *J Laryngol Otol*. 1991;105:436–8.
2. Menashe VD, Farrechic C, Miller M. Hypoventilation and corpulmonale due to chronic upper airway obstruction. *J Pediatr*. 1965;66:198–203.
3. Niedzielska G, Kotowski M, Niedzielski A. Assessment of pulmonary function and nasal flow in children with adenoid hypertrophy. *Int J Pediatr Otorhinolaryngol*. 2008;72:333–5.
4. Arens R, McDonough JM, Corbin AM, et al. Upper airway size analysis by magnetic resonance imaging of children with obstructive sleep apnea syndrome. *Am J Respir Crit Care Med*. 2003; 167:65–70.
5. Rosen CL, Larkin EK, Kirchner HL, et al. Prevalence and risk factors for sleep-disordered breathing in 8- to 11-year-old children: association with race and prematurity. *J Pediatr*. 2003; 142:383–9.
6. Tal A, Leiberman A, Margulis G, Sofer S. Ventricular dysfunction in children with obstructive sleep apnea: radionuclide assessment. *Pediatr Pulmonol*. 1988;4:139–43.
7. Marcus CL, Greene MG, Carroll JL. Blood pressure in children with obstructive sleep apnea. *Am J Respir Crit Care Med*. 1998;157:1098–103.
8. Gozal D. Sleep-disordered breathing and school performance in children. *Pediatrics*. 1998;102: 616–20.
9. O'Brien LM, Gozal D. Behavioral and neurocognitive implications of snoring and obstructive sleep apnea in children: facts and theory. *Paediatr Respir Rev*. 2002;3:3–9.
10. O'Brien LM, Mervis CB, Holbrook CR, Bruner JL, Klaus CJ, Rutherford J, et al. Neurobehavioral implications of habitual snoring in children. *Pediatrics*. 2004;114:44–9.
11. Kharb S, Yadav SPS, Singh H, Singh GP. Effect of adenotonsillectomy on arterial blood gases and acid-base balance. *Int J Pediatr Otorhinolaryngol*. 1998;43: 213–15.
12. Pires MG, Di Francesco RC, Grumach AS, Mello JF. Evaluation of inspiratory pressure in children with enlarged tonsils and adenoids. *Braz J Otorhinolaryngol*. 2005;71(5):598–601.
13. Mitchell RB, Kelly J. Quality of life after adenotonsillectomy for SDB in children. *Otolaryngol Head Neck Surg*. 2005;133(4):569–72.
14. Mitchell RB, Kelly J. Outcomes and quality of life following adenotonsillectomy for sleep-disordered breathing in children. *ORL J Otorhinolaryngol Relat Spec*. 2007;69(6):345–8.
15. Maurizi M, Paludetti G, Todisco T, Dottorini M, Grassi V. Pulmonary function studies in adenoid hypertrophy. *Int J Pediatr Otorhinolaryngol*. 1980; 2(3):243–50.
16. Di Martino E, Mlynski G, Mlynski B. Effect of adenoid hyperplasia on nasal airflow. *Laryngorhinology*. 1998;77:272–4.
17. Banzatto MG, Grumach AS, Mello JF Jr, Di Francesco RC. Adenotonsillectomy improves the strength of respiratory muscles in children with upper airway obstruction. *Int J Pediatr Otorhinolaryngol*. 2010;74(8):860–3.
18. Kavukcu S, Coskun S, Cevik N, Kuscü B, Akkoçlu A. The importance of pulmonary function tests in adenotonsillectomy indications. *Indian J Pediatr*. 1993;60:249–255.
19. Brodsky L. Tonsillitis, tonsillectomy and adenoidectomy. In: B.J. Bailey. *Head and Neck Surgery—Otolaryngology*. Philadelphia:Lippincott; 1993:833–47.
20. Fujioka M, Young LW, Girdany BR. Radiographic evaluation of adenoidal size in children: adenoidal- nasopharyngeal ratio. *Am J Roentgenol*. 1979;133:401–10.
21. Yernault JC, Englert M, Sergysels R, Decoster A. Upper airway stenosis: a physiologic study. *Am Rev Respir Dis*. 1973;108: 996–1000.
22. Yadav SP, Dodeja OP, Gupta KB, Chanda R. Pulmonary function tests in children with adenotonsillar hypertrophy. *Int J Pediatr Otorhinolaryngol*. 2003;67:121–5.
23. Hira HS, Singh H. Assessment of upper airway obstruction by pulmonary function testing. *J Assoc Physicians India*. 1994; 42:531–4.
24. Corren J. Allergic rhinitis and asthma, how important is the link? *J Allergy Clin Immunol*. 1997;99: S781–S6.
25. Geller M. Vias aéreas unidas, a atopia como processo sistêmico e contínuo. *Rev Bras Alerg Imunopatol*. 2002; 25:74–80.
26. Behlfelt K, Linderson-Aronson S, MacWilliam J, Neander P, Laage-Hellman J. Cranio-facial morphology in children with and without enlarged tonsils. *Eur J Orthod*. 1990;12(3):233–43.
27. Woodson GE, Robbins KT. Nasal obstruction and pulmonary function: The role of humidification. *Otolaryngol Head Neck Surg*. 1985;93:505–11.
28. Kendall FP, McCreary EK, Provance PG. *Muscles, Testing and Function with Posture and Pain*, 4th ed. Baltimore:Lippincott Williams & Wilkins; 2005.

- 29.** Blum DJ, Neel HB. Current thinking on tonsillectomy and adenoidectomy. *Comp Ther.* 1983; 9: 48–56.
- 30.** Doğruer ZN, Ünal M, Eskandari G, Pata YS, Akbaş Y, Cevik T, Cimen MY Malondialdehyde and antioxidant enzymes in children with obstructive adenotonsillar hypertrophy. *Clin Biochem.* 2004; 37:718–21.
- 31.** Ünal M, Öztürk C, Görür K. Effect of tonsillectomy on serum concentrations of interleukins and TNF- α in patients with chronic tonsillitis. *ORL J Otorhinolaryngol Relat Spec* 2002;64:254–6.
- 32.** Boyd JH, Petrof BJ, Hamid Q, Fraser R, Kimoff RJ. Upper airway muscle inflammation and denervation changes in obstructive sleep apnea. *Am J Respir Crit Care Med* 2004;170:541–6.
- 33.** Tauman R, Ivanenko A, O'Brien LM, Gozal D. Plasma C-reactive protein levels among children with sleep-disordered breathing. *Pediatrics*2004;113:564-9.
- 34.** Goldbart AD, Veling MC, Goldman JL, Li RC, Brittian KR, Gozal D. Glucocorticoid receptor subunit expression in adenotonsillar tissue of children with obstructive sleep apnea. *Pediatr Res.* 2005;57:232–6.
- 35.** Goldbart AD, Goldman GL, Li RC, Brittian KR, Tauman R, Gozal D. Differential expression of cysteinyl leukotriene receptors 1 and 2 in tonsils of children with obstructive sleep apnea and recurrent infection. *Chest.* 2004;126:13–18.
- 36.** Goldbart AD, Goldman JL, Veling MC, Gozal D. Leukotriene modifier therapy for mild sleep-disordered breathing in children. *Am J Respir Crit Care Med.* 2005;172:364–70.
- 37.** Kheirandish L, Goldbart AD, Gozal D. Intranasal Steroids and Oral Leukotriene Modifier Therapy in Residual Sleep-Disordered Breathing After Tonsillectomy and Adenoidectomy in Children *Pediatrics* 2006;117:e61–e6.